



Oral Vs Parenteral Iron Therapy among Women in Post Partum Period

Mamoona Suleman¹, Aneeza Akram² and Jawariya Arooj³

¹WMO in THQ Duniyapur

Email: mamoona2020@gmail.com

²Women medical officer

Email address: aneezaakram329@yahoo.com

³WMO in BHU Gujranwala

Email: jawariyaarooj97@gmail.com

Abstract

Objective: To determine efficacy of oral iron therapy as compared to parenteral iron therapy.

Study design and duration: This is a prospective study of observational type. Study consists on the duration of 5 months.

Setting: This study was conducted in Bahawal Victoria Hospital Bahawalpur.

Patients and Methods: Anemic patients having Hb less than 10g/dl were included in this study. Two groups were formed group-A and group-B. Group-A was given oral iron therapy while group-B was given Intravenous iron therapy and outcome was noted after 30 days and 60 days in the form of hemoglobin level, hematocrit and MCV values. All data collected was analyzed using SPSS and Microsoft office version 2016. Results were calculated in the form of frequencies and percentages and averages. Tables and Graphs used to present data. Consent was taken from ethical committee of the study hospital and consent was also taken from all cases in study group.

Results: There were total 70 cases in study group selected via randomized controlled trials. There were two groups containing 35 in each group. In group-A after 60 days of iron therapy via oral route HB value was 10.7 g/dl and after 60 days 11.8g/dl with the change of 22.4% in HB level. In Group-B after 60 days of intravenous iron therapy HB level was 12.9 g/dl with improvement of 38.8% in HB level. P- Value was 0.05. Age range of cases was 15-35 years of age with mean age of 31.4 years. Baseline characteristics of cases in group-A were age 33±4 years, BMI 23±5, HB level was 9.2±1 g/dl, Hematocrit (HCT) was 29.5±3 and MCV was 72±0.6. In group B mean age was 34±3, BMI 21±2, HB level 9.2±0.5, Hct was 28.6±3.4, MCV was 70±4.

Conclusion: Parenteral route of iron therapy is more effective in post partum period in women than oral route with more improvement in blood hemoglobin, hematocrit and MCV value.

Key Words: iron therapy, Oral route, Parenteral route, Hemoglobin, hematocrit, post partum period

Introduction

Women in post partum period usually develop anemia due to blood loss and in 30% cases Hb

level is less than 10g/dl. Anemia is defined as decreased in hemoglobin concentration and Hct as well. Severe anemia with Hb less than 8g/dl is found in 10% cases in post partum period. In pregnancy WHO defines anemia as Hb level less than 11g/dl in first and 3rd trimester and less than 10.5g/dl in 2nd trimester. It is a very common problem worldwide. Iron deficiency anemia in pregnancy is very common. During pregnancy maternal red cell mass is increased and fetoplacental demand for iron is increased as well leading to development of iron deficiency anemia. Iron deficiency during pregnancy and blood loss during delivery cause severe anemia in women and increasing demand for iron therapy. Iron therapy can be given via oral as well as parenteral route. Oral route is not much effective as compared to intravenous route due to absorption factor through intestine. Intravenous iron therapy is very effective in increasing hemoglobin and Hct level so correcting anemia in one to two months.

Patients and Methods


This is a prospective study in which effect of iron therapy is observed in two groups of patients. This study was started in January 2019 and completed in May 2019 consisting on total duration of 5 months. This study was conducted in a tertiary care hospital Bahawal Victoria Hospital located in Bahawalpur, A city of Pakistan. Anemic patients having Hb less than 10g/dl were included in this study. Two groups were formed group-A and group-B. Group-A was given oral iron therapy while group-B was given Intravenous iron therapy and outcome was noted after 30 days and 60 days in the form of hemoglobin level, hematocrit and MCV values. All data collected was analyzed using SPSS and Microsoft office version 2016. Results were calculated in the form of frequencies and percentages and averages. Tables and Graphs used to present data. Consent was taken from ethical committee of the study hospital and consent was also taken from all cases in study group.



Results

There were total 70 cases in study group selected via randomized controlled trials. There were two groups containing 35 in each group. In group-A after 60 days of iron therapy via oral route HB value was 10.7 g/dl and after 60 days 11.8g/dl with the change of 22.4% in HB level. In Group-B after 60 days of intravenous iron therapy HB level was 12.9 g/dl with improvement of 38.8% in HB level. P- Value was 0.05. Age range of cases was 15-35 years of age with mean age of 31.4 years. This is a prospective study in which effect of iron therapy is observed in two groups of patients. This study was started in January 2019 and completed in May 2019 consisting on total duration of 5 months. This

study was conducted in a tertiary care hospital Bahawal Victoria Hospital located in Bahawalpur, A city of Pakistan. Anemic patients having Hb less than 10g/dl were included in this study. Two groups were formed group-A and group-B. Group-A was given oral iron therapy while group-B was given Intravenous iron therapy and outcome was noted after 30 days and 60 days in the form of hemoglobin level, hematocrit and MCV values. Baseline characteristics of cases in group-A were age 33 ± 4 years, BMI 23 ± 5 , HB level was 9.2 ± 1 g/dl, Hematocrit (HCT) was 29.5 ± 3 and MCV was 72 ± 0.6 . In group B mean age was 34 ± 3 , BMI 21 ± 2 , HB level 9.2 ± 0.5 , Hct was 28.6 ± 3.4 , MCV was 70 ± 4 .



(Figure-1) Improvement in Hemoglobin via oral vs parenteral route

DISCUSSION

In women during pregnancy and after delivery anemia develops due to iron deficiency. This anemia needs to be corrected either in the form of oral iron therapy or intravenous therapy. Oral iron is not completely absorbed and many factors decrease its absorption so results are not much satisfactory. Iron given via intravenous route is much effective as 100% bioavailability and its rapid response corrects anemia within days. Usually iron is available in the form of tablets and injection ampoules which are mixed in fluid and given slowly intravenously. Intravenous iron may cause reaction so initially it is given very slowly to check any reaction. Women in post partum period usually develop anemia due to blood loss and in 30% cases Hb level is less than 10g/dl. Anemia is defined as decreased in hemoglobin concentration and Hct as well. There were total 70 cases in study

group selected via randomized controlled trials. There were two groups containing 35 in each group. In group-A after 60 days of iron therapy via oral route HB value was 10.7 g/dl and after 60 days 11.8g/dl with the change of 22.4% in HB level. In Group-B after 60 days of intravenous iron therapy HB level was 12.9 g/dl with improvement of 38.8% in HB level. P- Value was 0.05. Age range of cases was 15-35 years of age with mean age of 31.4 years. Severe anemia with Hb less than 8g/dl is found in 10% cases in post partum period. In pregnancy WHO defines anemia as Hb level less than 11g/dl in first and 3rd trimester and less than 10.5g/dl in 2nd trimester. This is a prospective study in which effect of iron therapy is observed in two groups of patients. This study was started in January 2019 and completed in May 2019 consisting on total duration of 5 months. This study was conducted in a tertiary care hospital Bahawal



Victoria Hospital located in Bahawalpur, A city of Pakistan. Anemic patients having Hb less than 10g/dl were included in this study. Two groups were formed group-A and group-B. It is a very common problem worldwide. Iron deficiency anemia in pregnancy is very common. During pregnancy maternal red cell mass is increased and fetoplacental demand for iron is increased as well leading to development of iron deficiency anemia. . All data collected was analyzed using SPSS and Microsoft office version 2016. Results were calculated in the form of frequencies and percentages and averages. Tables and Graphs used to present data. Consent was taken from ethical

committee of the study hospital and consent was also taken from all cases in study group. Iron deficiency during pregnancy and blood loss during delivery cause severe anemia in women and increasing demand for iron therapy. Iron therapy can be given via oral as well as parenral route.

Conclusion

Iron given intravenously corrects anemia rapidly as compared to orally given. It improves hemoglobin, hematocrit and MCV effectively. Oral iron therapy is less effective due to non-compliance, less bioavailability and more side effects.

REFERENCES

1. Wojtyła C, Biliński P, Paprzycki P, Warzocha K. **Haematological parameters in postpartum women and their babies in Poland – comparison of urban and rural areas.** Annals of Agricultural and Environmental Medicine 2011, Vol 18, No 2, 380-385
2. Bhagwan D, Kumar A, Rao CR, Kamath A. **Prevalence of Anaemia among Postnatal Mothers in Coastal Karnataka.** Journal of Clinical and Diagnostic Research. 2016 Jan, Vol-10(1).
3. Baker W. **Iron deficiency in pregnancy, obstetrics and gynecology.** Hematol Oncol Clin North Am 2000; 14:1061–77.
4. Bashiri A, Burstein E, Sheiner E, Mazor M. **Anemia during pregnancy and treatment with intravenous iron: Review of the literature.** European Journal of Obstetrics, Gynecology, & Reproductive Biology. 2003; 110:2-7.
5. Becuzzi N, Zimmermann R, Alexander Krafft A. **Long-Term Efficacy of Postpartum Intravenous Iron Therapy.** BioMed Research International Vol 2014, 5 pages.
6. Trinh LTT, Dibley M. **Anaemia in pregnant, postpartum and non-pregnant women in Lak district, Daklak province of Vietnam.** Asia Pac J Clin Nutr, 2007; 16 (2):310-315.
7. Rathod S, Samal SK, Mahapatra PC, S. **Ferric carboxymaltose: A revolution in the treatment of postpartum anemia in Indian women.** Int J Appl Basic Med Res. 2015 Jan-Apr; 5(1): 25–30.
8. Baig-Ansari N, Badruddin SH, Karmaliani R, Harris H, Jehan I, Pasha O, Moss N, McClure EM, Goldenberg. **Anemia prevalence and risk factors in pregnant women in an urban area of Pakistan.** Food Nutr Bull. 2008 June; 29(2): 132–139.
9. Bodnar LM, Scanlon KS, Freedman DS, Siega-Riz AM, Cogswell ME. **High prevalence of postpartum anemia among low-income women in the united states.** American Journal of Obstetrics & Gynecology. 2001; 185:438-443.
10. World Health Organization. **Reduction of maternal mortality. A joint WHO/UNFPA/UNICEF/World bank statement.** Geneva: WHO; 1999.
11. Breyman C. **Treatment of iron deficiency anaemia in pregnancy and postpartum with special focus on intravenous iron sucrose complex.** Journal of the Medical Association of Thailand. 2005; 88:S108-9.
12. Rienold C, Dalenius K, Smith B, Brindley P, Grummer-Strawn L. **Pregnancy nutrition surveillance 2007 report.** Atlanta: U.S. Department of Health and Human Services, Center for Disease Control and Prevention; 2009.
13. Zainur RZ, Loh KY. **Postpartum Morbidity - What We Can Do.** Med J Malaysia, December 2006, Vol 61, No. 5.
14. Beard JL, Hendricks MK, Perez EM, et al. **Maternal iron deficiency anemia affects postpartum emotions and cognition.** J Nutr. 2005; 135:267-272.



15. Rakesh PS, Gopichandran V, Jamkhandi D, Manjunath K, George K, Prasad J. **Determinants of postpartum anemia among women from a rural population in southern India.** International Journal of Women's Health 2014;6 395–400.
16. Perez EM, Hendricks MK, Beard JL, et al. **Mother-infant interactions and infant development are altered by maternal iron deficiency anemia.** J Nutr. 2005; 135:850-855.
17. Perello MF, Coloma JL, Masoller N, Esteve J, Palacio M. **Intravenous ferrous sucrose versus placebo in addition to oral iron therapy for the treatment of severe postpartum anaemia; a randomized controlled trial.** BJOG, 2014; 121: 706-713.
18. Breyman C, Gliga F, Bejenariu C, Strizhova N. **Comparative efficacy and safety of intravenous ferric carboxymaltose in the treatment of postpartum iron deficiency anemia.** International Journal of Gynaecology & Obstetrics. 2008; 101:67-73.
19. Perewunsky G, Huch R, Huch A, et al. **Parenteral iron therapy in obstetrics: 8 years experience with iron– sucrose complex.** Br J Nutr. 2002; 88:3–10.
20. Bhandal N, Russell R. **Intravenous versus oral iron therapy for postpartum anaemia.** BJOG. 2006; 113:1248–52.
21. Broche DE, Gay C, Armand-Branger S, et al. **Severe anaemia in the immediate post-partum period. Clinical practice and value of intravenous iron.** Eur J Obstet Gynecol Reprod Biol. 2005; 123(2):S21–7.
22. Geeta J, Urmila P, Jha SK. **Intravenous Iron in Postpartum Anemia.** The Journal of Obstetrics and Gynecology of India (January–February 2013) 63(1):45– 48.